

Entropy Corrections for Schwarzschild and Reissner-Nordström Black Holes

M. M. Akbar^b, Saurya Das[#] *

^b
Department of Applied Mathematics and Theoretical Physics,
Centre for Mathematical Sciences,
University of Cambridge,
Wilberforce Road,
Cambridge CB3 0WA, U.K.
Email: M.M.Akbar@damtp.cam.ac.uk

[#]
Department of Mathematics and Statistics,
University of New Brunswick,
Fredericton, New Brunswick E3B 5A3, CANADA
Email: saurya@math.unb.ca

Schwarzschild black hole being thermodynamically unstable, corrections to its entropy due to small thermal fluctuations cannot be computed. However, a thermodynamically stable Schwarzschild solution can be obtained within a cavity of any finite radius by immersing it in an isothermal bath. For these boundary conditions, classically there are either two black hole solutions or no solution. In the former case, the larger mass solution has a positive specific heat and hence is locally thermodynamically stable. We find that the entropy of this black hole, including first order fluctuation corrections is given by: $S = S_{BH} - \ln[\frac{3}{R}(S_{BH}/4\pi)^{1/2} - 2]^{-1} + (1/2)\ln(4\pi)$, where $S_{BH} = A/4$ is its Bekenstein-Hawking entropy and R is the radius of the cavity. We extend our results to four dimensional Reissner-Nordström black holes, for which the corresponding expression is: $S = S_{BH} - \frac{1}{2}\ln[(S_{BH}/\pi R^2)(3S_{BH}/\pi R^2 - 2\sqrt{S_{BH}/\pi R^2 - \alpha^2}) (\sqrt{S_{BH}/\pi R^2 - \alpha^2})/(S_{BH}/\pi R^2 - \alpha^2)^2]^{-1} + (1/2)\ln(4\pi)$. Finally, we generalise the stability analysis to Reissner-Nordström black holes in arbitrary spacetime dimensions, and compute their leading order entropy corrections. In contrast to previously studied examples, we find that the entropy corrections in these cases have a different character.

I. INTRODUCTION

It is well-known that the specific heat of a Schwarzschild black hole is negative:

$$C = \frac{dM}{dT} = -8\pi M^2 < 0. \quad (1)$$

Thus it is thermodynamically unstable, and corrections to thermodynamic quantities of this black hole, e.g. entropy and temperature, are not well-defined. This is seen when one considers for instance the corrected entropy of any thermodynamic system (including black holes) due to small thermal fluctuations of the system, which is given by [1,2]:

$$\begin{aligned} S &= S_0 - \frac{1}{2}\ln(<E^2> - <E>^2) + \dots \\ &= S_0 - \frac{1}{2}\ln(CT^2) + \dots \end{aligned} \quad (2)$$

When relation (2) is applied to black holes, one substitutes $S_0 \rightarrow S_{BH}$, the Bekenstein-Hawking entropy of the

black hole, and calculates C for the particular black hole under consideration. Note that in using (2) in the context of black holes, one is simply assuming that the black hole behaves as an ordinary thermodynamic system, following usual laws of thermodynamics. Then starting from a continuum partition function and performing an inverse Laplace transform gives the density of states, whose logarithm yields Eq.(2) above. In addition, if one assumes that the black hole (more precisely, its horizon) is built up of more fundamental entities, such as quanta of area, or that observables such as horizon area and mass are quantised in a certain way, then as shown in refs. [3] and [4], there could be additional terms in (2).

It may also be noted that there could be other corrections to black hole entropy due to a variety of other sources. For instance, quantum fluctuations of matter fields in black hole backgrounds, as well as fluctuations of spacetime geometry itself in the canonical quantum gravity framework give rise to corrections which are also logarithmic in nature. Details of these calculations can be found in [5–23]. Since our analysis is not tied to any specific model of quantum gravity, it remains valid as

*Address from August 2003: Dept. of Physics, University of Lethbridge, 4401 University Drive, Lethbridge, Alberta T1K 3M4, CANADA. Email: saurya.das@uleth.ca

long as the black hole is large. On the other hand, the other corrections may become important as the size of the black hole approaches the Planck scale.

One sees that the formula breaks down for $C < 0$. However, as was first shown by York [24], a positive specific heat solution can be obtained by considering Schwarzschild black holes within a *finite* S^2 cavity immersed in an isothermal bath. In fact for such canonical boundary conditions there are either two or no solutions depending on whether the temperature of the bath is above or below a minimum value. The larger mass of the solutions has a positive heat and the other has a negative specific heat. When one takes the cavity radius to infinity keeping the wall-temperature of the cavity fixed (i.e., not changing the temperature of the bath) the positive specific heat solution disappears and only the negative specific solution survives giving the known black hole solution. This holds for higher dimensions as well [25]. In this article, we consider the positive specific heat solution and compute leading order corrections to its entropy. As we will see the logarithmic correction term due to small thermal fluctuation is of a different nature unlike in several other cases studied before [5–23]. In this case, the mass and specific heat of the black hole are functions of the temperature *and* cavity radius, and the entropy correction is more interesting and non-trivial, as we will see below. The case of charged black hole, i.e., Reissner-Nordström black holes, within a cavity was first studied in [26] and we will find correction for their entropy as well for thermal fluctuation. See also [27] for related work on thermodynamics of Schwarzschild and Reissner-Nordström-AdS black holes. This paper is arranged in the following way. In the next section, we briefly review the thermodynamics of Schwarzschild black holes within a cavity and compute the entropy correction. In section (III), we extend our results to Reissner-Nordström black holes in arbitrary spacetime dimensions. Finally, we end with a summary of our results and some further directions in sections (IV). We use units in which $G = c = \hbar = k_B = 1$.

II. SCHWARZSCHILD BLACK HOLE IN A BOX: ENTROPY CORRECTIONS

Let us first consider a Schwarzschild black hole of mass M . Wick rotating the Lorentzian time $t \rightarrow i\tau$ one obtains

$$ds^2 = \left(1 - \frac{2M}{r}\right) d\tau^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2 d\Omega_2^2, \quad (3)$$

where $d\Omega_2^2$ is the metric on unit S^2 . The metric (3) is singular at $r = 2M$. The singularity can be removed by periodically identifying τ with a period $8\pi M$. Hence the

resulting metric is complete and regular for $2M \leq r < \infty$. The fixed point set of the Killing vector $\partial/\partial\tau$ is a regular bolt of the metric [28]. The inverse of the periodicity of the τ coordinate gives the temperature of the hole measured at infinity.

A. Isothermal Cavity

Next, consider a spherical S^2 cavity of radius R concentric with the black hole horizon. The black hole temperature measured at infinity is shifted by $(g_{00})^{-\frac{1}{2}}$, as one moves towards the black hole. The local temperature of the S^2 cavity is therefore given by

$$\beta = T^{-1} = 8\pi M \sqrt{1 - \frac{2M}{R}}. \quad (4)$$

Now we ask the reverse question. Given a spherical cavity for radius R held at a temperature β^{-1} , what are the black hole solutions? In other words what are possible solutions for M ? This was first considered by York [24] for four dimensions. Recently this has been discussed for higher dimensions in [25]. For the purpose of exposition we deal with the case of four dimensions here and discuss generalisation to arbitrary dimensions in the next section. It is not difficult to see that there are in fact two black hole solutions or no solutions by considering Eq.(4) which can be written in the following form :

$$M^3 - \frac{1}{2}RM^2 + \frac{R\beta^2}{128\pi^2} = 0. \quad (5)$$

This is a cubic equation in M and always admits a negative root and a pair of positive or complex roots depending on the coefficients. The coefficients are determined by the boundary variables R and β . Defining new variables $x \equiv 2M/R$ and $\sigma = \beta/4\pi R$, Eq.(5) can be cast in the following simple form:

$$x^3 - x^2 + \sigma^2 = 0. \quad (6)$$

When positive solutions exist, larger and smaller solutions of x give larger and smaller hole respectively. Note that the positive roots take values from zero to one (see figure 1). It can be shown that two positive roots occurs if and only if [24]

$$\sigma \leq \frac{2}{\sqrt{27}}. \quad (7)$$

Thus one requires the temperature T of the bath to be

$$T \geq \frac{\sqrt{27}}{8\pi} \frac{1}{R}. \quad (8)$$

At the equality of (7) or (8) the two roots are degenerate. The larger solution therefore takes value within the closed interval $[\frac{2}{3}, 1]$. For completeness we mention here the explicit solutions of (6) here. The exact solutions are:

$$x_1 = \frac{1}{3} \left[1 - 2 \cos \left(\frac{\alpha}{3} + \frac{\pi}{3} \right) \right] , \quad (9)$$

$$x_2 = \frac{1}{3} \left[1 + 2 \cos \left(\frac{\alpha}{3} \right) \right] , \quad (10)$$

which translate in terms of black hole mass as:

$$M_1 = \frac{R}{6} \left[1 - 2 \cos \left(\frac{\alpha}{3} + \frac{\pi}{3} \right) \right] , \quad (11)$$

$$M_2 = \frac{R}{6} \left[1 + 2 \cos \left(\frac{\alpha}{3} \right) \right] , \quad (12)$$

where

$$\cos \alpha = 1 - \frac{27}{2} \sigma^2 . \quad (13)$$

Evidently at the equality of (7), $\alpha = \pi$, the solutions coincide, and

$$x = \frac{2}{3} \Leftrightarrow M = \frac{R}{3} . \quad (14)$$

By calculating the Euclidean action of the solution it was shown in [24] that the entropy of the black hole is unaffected by the presence of the box and is given by the usual relationship

$$S_{BH} = 4\pi M^2 . \quad (15)$$

The specific heat is defined as

$$C_R = T \left(\frac{\partial S_{BH}}{\partial T} \right)_R , \quad (16)$$

From (4) and (15), one obtains:

$$C = 8\pi M^2 \left(\frac{R - 2M}{3M - R} \right) = 4\pi R^2 x^2 \left(\frac{1 - x}{3x - 2} \right) . \quad (17)$$

From its form it is apparent that the specific heat is positive in the range

$$\frac{2}{3} \leq x \leq 1 \Leftrightarrow 2M \leq R \leq 3M , \quad (18)$$

which is possible only for the larger hole (M_2 above). This can be also observed from Figure 1. For constant value of R , Figure 1 is a relation between (inverse of) temperature and the masses of the two Schwarzschild black holes. The mass of the larger solution increases with increasing temperature. For the smaller hole it is just the opposite. Also note that in the $R \rightarrow \infty$ limit, the larger mass solutions do not exist meaningfully (it fills in the cavity) irrespective of the temperature. Only the smaller solution x_1 exists with $M = \frac{\beta}{8\pi}$ since $x_1 \equiv \frac{2M}{R} = \frac{\beta}{4\pi R}$ in this limit.




FIG. 1. For every value of $\sigma^2 \equiv 1/(4\pi RT)^2 < \frac{4}{27}$ there are two values of x , i.e., two black holes. For the larger black hole $x \in [\frac{2}{3}, 1]$.

B. Logarithmic Correction

Substituting (17) and (4) in (2), we obtain the leading corrections to the entropy of the larger black hole:

$$\begin{aligned} \mathcal{S} &= S_{BH} - \frac{1}{2} \ln((3x - 2)^{-1}) + \frac{1}{2} \ln(4\pi) \dots \\ &= S_{BH} - \frac{1}{2} \ln \left[\frac{3}{R} \left(\frac{S_{BH}}{4\pi} \right)^{1/2} - 2 \right]^{-1} + \frac{1}{2} \ln(4\pi) + \dots \end{aligned} \quad (19)$$

Note that the first correction term is negative. This is because, since $x \in [2/3, 1]$, the argument of the logarithm is always greater than unity. Also, note that the corrections here is of a somewhat different type than those found in the different applications of (2). First, the correction depends on the value of x which is a function of the product RT . When $RT \rightarrow \frac{\sqrt{27}}{8\pi}$ the correction is enormous irrespective of the value of R alone (the horizon radius is $\frac{2R}{3}$). As one increases the temperature, keeping the radius of the cavity fixed, the infilling black hole becomes larger and fills in the whole cavity at infinite temperature. However, the correction term decreases with increasing temperature and still remains finite and non-zero approaching $\ln(4\pi)$ at the infinite temperature limit. For small cavity radius the correction term is therefore given by the first term and is always appreciable and overpowers S_{BH} if the temperature is not high enough (signaling breakdown of formalism). For large cavity radius it is more appreciable at lower temperatures and becomes negligible, albeit non-zero, at high temperature. Again, if T is near $\frac{\sqrt{27}}{8\pi R}$ it overpowers S_{BH} even if the latter is large (again signaling a breakdown).

Another way to see the different nature of the correction terms is to note the coefficient of the correction term. The coefficient in front of either of the two terms (which can be combined together) is $1/2$, in contrast to BTZ of Schwarzschild anti-de Sitter black holes for example, where the coefficients are $-3/2$ and $-d/2(d-2)$ respectively [1]. However note that this $1/2$ is not in the same

footing as the coefficients in those cases as the correction term here is not of the type $-k \ln(S_{BH})$. Rather, if we are to write it in terms of S_{BH} it takes the form given by the second equation of (19) which involves the cavity radius as well.

C. Higher Dimensions

Higher dimensional Schwarzschild black holes and their thermodynamics within a finite isothermal cavity have been studied in [25]. For d -dimensions the inverse temperature at the $(d-2)$ -dimensional boundary is given by:

$$\beta = T^{-1} = \frac{4\pi x R}{d-3} \sqrt{1-x^{d-3}} \equiv \frac{4\pi R \sigma}{d-3}, \quad (20)$$

from which one obtains:

$$x^{d-1} - x^2 + \sigma^2 = 0. \quad (21)$$

This equation cannot be solved using ordinary algebraic methods. However, analytic solution is still possible [25]. This would be needed if one requires the entropy correction as a function of cavity radius and temperature. However, as in the above case of four dimensions the general properties of the corrections to entropy can be studied without requiring the explicit solutions by studying how the correction term varies with x . (Again note that x takes value within the interval $[0, 1]$.) This will be done in the next section as a special case of d -dimensional Reissner-Nordström. As we will see there entropy corrections to d -dimensional Reissner-Nordström (and hence Schwarzschild) black holes can be exactly evaluated.

III. REISSNER-NORDSTRÖM BLACK HOLES

We now extend the above analysis for the Reissner-Nordström black holes which have a charge. This means we would have to specify suitable electrostatic potential at the boundary. We first discuss the case of four dimensions.

A. Four dimensions

The Euclideanised Reissner-Nordström black hole metric is:

$$ds^2 = \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2}\right) d\tau^2 + \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2}\right)^{-1} dr^2 + r^2 d\Omega_2^2, \quad (22)$$

with inverse temperature at boundary [26]:

$$\begin{aligned} \beta = T^{-1} &= 4\pi r_+ \left(1 - \frac{Q^2}{r_+^2}\right)^{-1} \left(1 - \frac{r_+}{R}\right)^{1/2} \left(1 - \frac{Q^2}{r_+ R}\right)^{1/2} \\ &= \frac{4\pi R x^{5/2}}{x^2 - \alpha^2} \sqrt{(1-x)(x-\alpha^2)}, \end{aligned} \quad (23)$$

where $r_+ = M + \sqrt{M^2 - Q^2}$ is the radius of the outer horizon and

$$\alpha = \frac{Q}{R}.$$

The counterpart of Eq.(5) is:

$$(1 - \Phi^2)x^3 - x^2 + (1 - \Phi^2)^2 \sigma^2 = 0, \quad (24)$$

where x and σ has identical definitions in terms of r_+ and β as before. Here, Φ is the potential difference between r_+ and R , suitably red-shifted:

$$\begin{aligned} \Phi &= \frac{Q}{r_+} \left(1 - \frac{r_+}{R}\right)^{1/2} \left(1 - \frac{Q^2}{R r_+}\right)^{-1/2} \\ &= \alpha \sqrt{\frac{1-x}{x^2 - \alpha^2 x}}, \end{aligned} \quad (25)$$

Now, the two positive solutions of x are:

$$x_1 = \frac{1}{3(1-\Phi^2)} \left(1 - 2 \cos\left(\frac{\alpha}{3} + \frac{\pi}{3}\right)\right) \quad (26)$$

$$x_2 = \frac{1}{3(1-\Phi^2)} \left(1 + 2 \cos\frac{\alpha}{3}\right) \quad (27)$$

$$\cos \alpha = 1 - \frac{27}{2} \sigma^2 (1-\Phi)^4, \quad (28)$$

of which only the second one is thermodynamically stable. The entropy is, in this case:

$$S_{BH} = \pi r_+^2 = \pi R^2 x^2. \quad (29)$$

(where we have dropped the subscript 2 from x_2). The specific heat at constant Φ and R can now be computed by using the relation:

$$C_{\Phi,R} = \left(\frac{\partial E}{\partial T}\right)_{\Phi,R} = - \left(\frac{\partial S / \partial x}{\partial \ln \sigma / \partial x}\right)_{\Phi,R} \quad (30)$$

with σ obtained from Eq.(24). This yields:

$$C_{\Phi,R} = \frac{4\pi R^2 x^3 (1-x)}{3x^2 - 2x - \alpha^2}, \quad (31)$$

where now the specific heat is non-negative in the range:

$$r_+ \leq R \leq \frac{3}{2} r_+ - \frac{1}{2} \frac{Q^2}{r_+}, \quad (32)$$

Equivalently:

$$x \leq 1, \quad (33)$$

$$3x^2 - 2x - \alpha^2 \geq 0. \quad (34)$$

In the above range, the corrected entropy assumes the form:

$$\mathcal{S} = S_{BH} - \frac{1}{2} \ln \left(\frac{x^2(3x^2 - 2x - \alpha^2)(x - \alpha^2)}{(x^2 - \alpha^2)^2} \right)^{-1} + \frac{1}{2} \ln(4\pi) + \dots \quad (35)$$

This can be expressed in terms of the Bekenstein-Hawking entropy as:

$$\mathcal{S} = S_{BH} - \frac{1}{2} \ln \left(\frac{\left(\frac{3S_{BH}}{\pi R^2} - 2\sqrt{\frac{S_{BH}}{\pi R^2}} - \alpha^2 \right) \left(\sqrt{\frac{S_{BH}}{\pi R^2}} - \alpha^2 \right)}{\left(\frac{\pi R^2}{S_{BH}} \right) \left(\frac{S_{BH}}{\pi R^2} - \alpha^2 \right)^2} \right)^{-1} + \frac{1}{2} \ln(4\pi) + \dots \quad (36)$$

As expected, the above reduces to Eq.(19) for $\alpha = 0$.

B. Higher dimensions

For the d -dimensional Reissner-Nordström black hole the Euclidean metric is given by

$$ds^2 = \left(1 - \frac{16\pi M}{(d-2)\Omega_{d-2}r^{d-3}} + \frac{16\pi Q^2}{(d-2)(d-3)r^{2(d-3)}} \right) d\tau^2 + \left(1 - \frac{16\pi M}{(d-2)\Omega_{d-2}r^{d-3}} + \frac{16\pi Q^2}{(d-2)(d-3)r^{2(d-3)}} \right)^{-1} dr^2 + r^2 d\Omega_{d-2}^2, \quad (37)$$

where $d\Omega_{d-2}^2$ is a unit S^{d-2} sphere with the standard round metric on it. The inverse temperature is given by:

$$\beta = T^{-1} = \frac{4\pi r_+}{d-3} \left(1 - \frac{2Q^2}{(d-2)(d-3)r_+^{2(d-3)}} \right)^{-1} \times \left(1 - \left(\frac{r_+}{R} \right)^{d-3} \right)^{\frac{1}{2}} \left(1 - \frac{2Q^2}{(d-2)(d-3)(r_+R)^{d-3}} \right)^{\frac{1}{2}} \quad (38)$$

$$= \frac{4\pi R x^{1+3(d-3)/2}}{d-3} \left(x^{2(d-3)} - \alpha^{2(d-3)} \right)^{-1} \times (1 - x^{d-3})^{1/2} \left(x^{d-3} - \alpha^{2(d-3)} \right)^{1/2},$$

where $x = r_+/R$ as before, and now α is defined as:

$$(\alpha R)^{2(d-3)} = \frac{2Q^2}{(d-2)(d-3)}.$$

Similarly, the the potential difference is:

$$\Phi = \frac{Q}{r_+^{d-3}} \sqrt{\frac{1 - (r_+/R)^{d-3}}{1 - 2Q^2/(d-2)(d-3)(r_+R)^{d-3}}} \quad (40)$$

$$= \sqrt{\frac{2}{(d-2)(d-3)}} \alpha^{d-3} \sqrt{\frac{1 - x^{d-3}}{x^{2(d-3)} - \alpha^{2(d-3)} x^{d-3}}} \quad (41)$$

and Eq.(24) is generalised to:

$$\left(1 - \frac{(d-2)(d-3)}{2} \Phi^2 \right) x^{d-1} - x^2 + \left(1 - \frac{(d-2)(d-3)}{2} \Phi^2 \right)^2 \sigma^2 = 0, \quad (42)$$

where σ has been defined as in Eq.(20). Using (30), the specific heat at constant R and Φ can now be computed as:

$$C_{\Phi,R} = \frac{(d-2)\Omega_{d-2}R^{d-2}x^{2d-5}(1-x^{d-3})}{(d-1)x^{2(d-3)} - 2x^{d-3} - (d-3)\alpha^{2(d-3)}}. \quad (43)$$

Stability is now guaranteed if the parameters lie in a range such that:

$$(d-1)x^{2(d-3)} - 2x^{d-3} - (d-3)\alpha^{2(d-3)} \geq 0. \quad (44)$$

Explicit evaluation of $x(M, Q)$ and $R(M, Q)$ from the above is complicated for higher dimensions though, because of the higher powers of x . Eq.(43), along with (39) now gives the corrected entropy as:

$$\mathcal{S} = S_{BH} - \frac{1}{2} \ln \left[x^{d-2} \left((d-1)x^{2(d-3)} - 2x^{d-3} - (d-3)\alpha^{2(d-3)} \right) \times \frac{x^{d-3} - \alpha^{2(d-3)}}{R^{d-4} (x^{2(d-3)} - \alpha^{2(d-3)})^2} \right]^{-1} + \frac{1}{2} \ln \left[\frac{(d-2)\Omega_{d-2}}{(4\pi)^2} \right] + \dots \quad (45)$$

which can be expressed in terms of S_{BH} as:

$$\mathcal{S} = S_{BH} - \ln \left[\left(\frac{4S_{BH}}{\Omega_{d-2}R^{d-2}} \right) \times \left\{ (d-1) \left(\frac{4S_{BH}}{\Omega_{d-2}R^{d-2}} \right)^{2(d-3)/(d-2)} - 2 \left(\frac{4S_{BH}}{\Omega_{d-2}R^{d-2}} \right)^{(d-3)/(d-2)} - (d-3)\alpha^{2(d-3)} \right\} \times \frac{\left\{ \left(\frac{4S_{BH}}{\Omega_{d-2}R^{d-2}} \right)^{(d-3)/(d-2)} - \alpha^{2(d-3)} \right\}}{\left\{ \left(\frac{4S_{BH}}{\Omega_{d-2}R^{d-2}} \right)^{2(d-3)/(d-2)} - \alpha^{2(d-3)} \right\}^2} \right]^{-1} + \frac{1}{2} \ln \left[\frac{(d-2)\Omega_{d-2}}{(4\pi)^2} \right] + \dots \quad (46)$$

For zero-charge, that is for d -dimensional Schwarzschild black holes, the corresponding expression is:

$$\mathcal{S} = S_{BH} - \frac{1}{2} \ln \left[\frac{(d-1) \left(\frac{4S_{BH}}{\Omega_{d-2}R^{d-2}} \right)^{(d-3)/(d-2)-2}}{R^{d-4} \left(\frac{4S_{BH}}{\Omega_{d-2}R^{d-2}} \right)^{(d-4)/(d-2)}} \right]^{-1} + \frac{1}{2} \ln \left[\frac{(d-2)\Omega_{d-2}}{(4\pi)^2} \right] + \dots \quad (47)$$

In each case, the entropy correction is different in nature than in previously studied examples.

IV. DISCUSSIONS

In this article, we have shown how leading order corrections due to thermal fluctuations can be obtained for Schwarzschild and Reissner-Nordström black holes within a finite cavity. With appropriate boundary data, within a cavity thermodynamically stable solutions are possible. Normally, such corrections cannot be computed for Schwarzschild owing to its negative specific heat. For Reissner-Nordström black holes, corrections of this type were computed in [1] for a very small range of parameters near extremality, and that too in a perturbative approach (with perturbation parameter Q/M). Here, the result is non-perturbative, and holds for any parameter range. Finally, we generalised the stability analysis to higher dimensional Reissner-Nordström black holes, and computed its leading order entropy corrections. We found that the leading logarithmic corrections come with a negative signature. This means that the corrected dimensionality of the Hilbert space of the black holes under consideration, which is $\mathcal{N} = \exp(\mathcal{S})$, is smaller than the uncorrected dimension. It would be interesting to explore implications of this result. It would also be interesting to do the generalisation to other types of black holes, e.g. Kerr-Newmann as well as black holes which are not asymptotically flat, e.g. those that are asymptotically anti-de Sitter or de-Sitter. The implications of the latter to AdS/CFT and the dS/CFT correspondence may turn out to be interesting. We hope to report on this elsewhere.

Acknowledgements SD thanks R. K. Bhaduri and P. Majumdar for useful correspondence and interesting discussions. This work is supported in part by the Natural Sciences and Engineering Research Council of Canada. MMA would like to thank the Overseas Research Scheme, the Cambridge Commonwealth Trust and DAMTP for financial support.

-
- [1] S. Das, P. Majumdar, R. K. Bhaduri, *Class. Quant. Grav.* **19** (2002) 2355 (hep-th/0111011).
 - [2] S. Das, in *Talk given at Canadian Association of Physics Annual Congress, Quebec City, June 2-5, 2002* (hep-th/0207072).
 - [3] G. Gour, A. J. M. Medved, *Class. Quant. Grav.* **20** (2003) 3307 (gr-qc/0305018).
 - [4] R. K. Bhaduri, M. N. Tran, S. Das, gr-qc/0312023.
 - [5] D. Fursaev, *Phys. Rev.* **D51** (1995) 5352 (hep-th/9412161).
 - [6] S. N. Solodukhin, *Phys. Rev.* **D57** (1998) 2410 (hep-th/9701106).
 - [7] R. B. Mann, S. N. Solodukhin, *Nucl. Phys.* **B523** (1998) 293 (hep-th/9709064); A. J. M. Medved, G. Kunstatter, *Phys. Rev.* **D60** (1999) 104029 (hep-th/9904070); A. J. M. Medved, G. Kunstatter, *Phys. Rev.* **D63** (2001) 104005 (hep-th/0009050).
 - [8] S. Nojiri, S. D. Odintsov, *Int. J. Mod. Phys.* **A 15** (2000) 989 (hep-th/9905089).
 - [9] R. K. Kaul and P. Majumdar, *Phys. Rev. Lett.* **84** (2000) 5255 (gr-qc/0002040).
 - [10] O. Obregon, M. Sabido, V. I. Tkach, *Gen. Rel. Grav.* **33** (2001) 913 (gr-qc/0003023).
 - [11] S. Carlip, *Class. Quant. Grav.* **17** (2000) 4175 (gr-qc/0005017).
 - [12] J. Jing, M-L Yan, *Phys. Rev.* **D63** (2001) 24003 (gr-qc/0005105).
 - [13] S. Das, R. K. Kaul and P. Majumdar, *Phys. Rev.* **D 63** (2001) 044019 (hep-th/0006211); P. Majumdar, *Pramana* **55** (2000) 511 (hep-th/0009008), hep-th/0011284, hep-th/0110198.
 - [14] D. Birmingham, S. Sen, *Phys. Rev.* **D63** (2001) 47501 (hep-th/0008051); D. Birmingham, I. Sachs, S. Sen, *Int. J. Mod. Phys.* **D10** (2001) 833 (hep-th/0102155).
 - [15] T. R. Govindarajan, R. K. Kaul, V. Suneeta, *Class. Quant. Grav.* **18** (2001) 2877 (gr-qc/0104010).
 - [16] K. S. Gupta, S. Sen, *Phys. Lett.* **526** (2002) 121 (hep-th/0112041), hep-th/0203094; K. S. Gupta, hep-th/0204137.
 - [17] S. A. Major, K. L. Setter, *Class. Quant. Grav.* **18** (2001) 5125 (gr-qc/0101031); *Class. Quant. Grav.* **18** (2001) 5293 (gr-qc/0108034).
 - [18] A. J. M. Medved, *Class. Quant. Grav.* **19** (2002) 2503 (hep-th/0201079) and gr-qc/0211004.
 - [19] K. S. Gupta, hep-th/0204137.
 - [20] A. J. M. Medved, *Class. Quant. Grav.* **20** (2003) 2147 (hep-th/0210017).
 - [21] G. Gour, *Phys. Rev.* **D66** (2002) 104022 (gr-qc/0210024).
 - [22] A. Chatterjee, P. Majumdar, gr-qc/0303030.
 - [23] S. Das, V. Husain, *Class. Quant. Grav.* **20** (2003) 4387 (hep-th/0303089).
 - [24] J. W. York, *Phys. Rev.* **D33** (1986) 2092.
 - [25] M. M. Akbar, G. W. Gibbons, *Class. Quant. Grav.* **20** (2003) 1787 (hep-th/0301026).
 - [26] H. W. Braden, J. D. Brown, B. F. Whiting and J. W. York, *Phys. Rev. D* **42** (1990) 3376.
 - [27] J. Louko, B. F. Whiting, *Phys. Rev.* **D51** (1995) 5583 (gr-qc/9411017); J. Louko, S. N. Winters-Hilt, *Phys. Rev.* **D54** (1996) 2647 (gr-qc/9602003).
 - [28] G. W. Gibbons and S. W. Hawking, *Commun. Math. Phys.* **66** (1979) 291.